

Research

Open Access

Outlining eicosanoid biosynthesis in the crustacean *Daphnia*

Lars-Henrik Heckmann*^{1,2}, Richard M Sibly¹, Martijn JTN Timmermans³ and Amanda Callaghan¹

Address: ¹University of Reading, School of Biological Sciences, Environmental Biology, PO Box 68, Reading, RG6 6BX, UK, ²University of Aarhus, National Environmental Research Institute, Department of Terrestrial Ecology, Vejløvej 25, PO Box 314, DK-8600, Silkeborg, Denmark and ³VU University Amsterdam, Institute of Ecological Science, Department of Animal Ecology, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands

Email: Lars-Henrik Heckmann* - lhh@dmu.dk; Richard M Sibly - r.m.sibly@reading.ac.uk; Martijn JTN Timmermans - martijn.timmermans@falw.vu.nl; Amanda Callaghan - a.callaghan@reading.ac.uk

* Corresponding author

Published: 14 July 2008

Received: 28 March 2008

Frontiers in Zoology 2008, 5:11 doi:10.1186/1742-9994-5-11

Accepted: 14 July 2008

This article is available from: <http://www.frontiersinzoology.com/content/5/1/11>

© 2008 Heckmann et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Eicosanoids are biologically active, oxygenated metabolites of three C20 polyunsaturated fatty acids. They act as signalling molecules within the autocrine or paracrine system in both vertebrates and invertebrates mainly functioning as important mediators in reproduction, the immune system and ion transport. The biosynthesis of eicosanoids has been intensively studied in mammals and it is known that they are synthesised from the fatty acid, arachidonic acid, through either the cyclooxygenase (COX) pathway; the lipoxygenase (LOX) pathway; or the cytochrome P450 epoxygenase pathway. However, little is still known about the synthesis and structure of the pathway in invertebrates.

Results: Here, we show transcriptomic evidence from *Daphnia magna* (Crustacea: Branchiopoda) together with a bioinformatic analysis of the *D. pulex* genome providing insight on the role of eicosanoids in these crustaceans as well as outlining a putative pathway of eicosanoid biosynthesis. *Daphnia* appear only to have one copy of the gene encoding the key enzyme COX, and phylogenetic analysis reveals that the predicted protein sequence of *Daphnia* COX clusters with other invertebrates. There is no current evidence of an epoxygenase pathway in *Daphnia*; however, LOX products are most certainly synthesised in daphnids.

Conclusion: We have outlined the structure of eicosanoid biosynthesis in *Daphnia*, a key genus in freshwater ecosystems. Improved knowledge of the function and synthesis of eicosanoids in *Daphnia* and other invertebrates could have important implications for several areas within ecology. This provisional overview of daphnid eicosanoid biosynthesis provides a guide on where to focus future research activities in this area.

Background

Eicosanoids are cell signalling molecules derived from fatty acids acquired in the diet. Eicosanoid is a general term for all biologically active, oxygenated metabolites of three C20 polyunsaturated fatty acids; 20:3 n-6, 20:4 n-6

and 20:5 n-3. They have an important role in the regulation of essential functions such as reproduction and the immune system. All mammalian eicosanoids derive from a common precursor, arachidonic acid (AA), which is converted into eicosanoids with different functions through

either the cyclooxygenase (COX) pathway (prostanoids: prostaglandins and thromboxane); the lipoxygenase (LOX) pathway (leukotrienes and lipoxins); or the cytochrome P450 epoxygenase pathway (epoxyeicosatrienoic acids) [1] (Fig. 1).

The COX enzyme exists in at least two isoforms in mammals, COX-1 and COX-2, which are inhibited by conventional non-steroidal anti-inflammatory drugs (NSAID) such as ibuprofen and aspirin, but only one COX isoform is generally present in invertebrates and lower vertebrates [2]. Mammalian COX-1 catalyses the generation of pros-




Figure 1

Overview of eicosanoid biosynthesis based on current knowledge from mammalian models. The three major pathways, cyclooxygenase (COX), lipoxygenase (LOX) and cytochrome P450 epoxygenase, are shown displaying major metabolites. Prostanoids cover prostaglandins (PG) and thromboxane (TX), while leukotrienes (LT) include LTs and lipoxins. PGD₂ and PGE₂ may be transformed into PGJ₂ and PGA₂ through either non-enzymatic rearrangement or dehydration, respectively. Abbreviations: EET, epoxyeicosatrienoic acids; HETE, hydroxyeicosatetraenoic acids; HPETE, hydroperoxyeicosatetraenoic acids. Diagram modified from Stanley [1] and KEGG [41].

taglandins (PG) involved in many basic physiological functions such as regulation of blood pressure, gastric mucosal protection, maintenance of homeostasis, and reproductive and nervous system function; while PGs metabolised by COX-2 are involved in inflammation, ovulation and mitogenesis [3,4]. COX-1 is a constitutive enzyme (i.e. it is constantly expressed) found in the cell membranes of most mammalian tissues, while COX-2 is an inducible enzyme, not found in all cell types, that is located in nuclear membranes [1]. In the mammalian LOX pathway AA is converted by different LOXs into hydroperoxyeicosatetraenoic acids (HPETE) that may be further metabolised into hydroxyeicosatetraenoic acids (HETE) by glutathione peroxidase. Leukotrienes are synthesised from 5-HPETE by 5-LOX, while 8-LOX, 12-LOX and 15-LOX catalyse the production of different lipoxins from 8-HPETE, 12-HPETE and 15-HPETE (Fig. 1).

Research on eicosanoids has mainly been mammalian-driven and has lately been aimed at designing NSAIDs that are COX-2 selective due to the potential negative side-effects of COX-1 inhibition, which may affect the gastrointestinal tract, heart and kidneys [3]. Eicosanoids act at both the extracellular and intracellular level by interacting with distinct transmembrane G-protein coupled receptors (extra- and intracellular) and nuclear peroxisomal proliferator-activated receptors (PPAR) [5]. Activated G-proteins may, depending on cell type, stimulate second messengers such as cyclic adenosine monophosphate (cAMP) and/or intracellular calcium release [5]. PPAR are transcription factors which also have a role in ligand binding (eicosanoids), so directly influencing the expression of target genes involved in e.g. controlling prenatal and postnatal development [6]. These examples emphasise the biological significance of eicosanoids.

Less is known about eicosanoids in non-mammalian species; however, during the last three decades considerable evidence has been gathered concerning their synthesis and action. Eicosanoids have now been identified in almost every major metazoan phyla including some plants [1]. There is general consensus that eicosanoids act as autocrine or paracrine signalers (also referred to as local hormones) in both vertebrates and invertebrates, where they mainly function as important mediators in reproduction, the immune system and ion transport [1]. It is clear from a number of reports that eicosanoid generation is subject to inhibition by NSAIDs in a wide range of invertebrates [1,2]. Moreover, a COX derived mechanism similar to the mammalian biosynthesis of PGs has been proposed in the coral *Plexaura homomalla* [7]. There is also evidence of a LOX derived pathway being present in invertebrates based on the work of Ragab and colleagues [8] on the primitive wingless insect, the firebrat *Thermobia domestica*; although little is known about the structure of the

pathway. Overall, invertebrate eicosanoid biosynthesis seems to have a simpler structure than its mammalian counterpart, as seen with the COX pathway [7], but also appears to be split into two instead of three pathways. There is currently no proof of an epoxygenase pathway being present in invertebrates [1].

There is little doubt that disruption of eicosanoid biosynthesis may upset many important physiological functions in both invertebrates and vertebrates, which can have serious consequences for both the individual and the population. Current evidence suggests that the main mode of action of the NSAID ibuprofen in *D. magna* relates to interruption of eicosanoid biosynthesis which reduces fecundity [9-12]. Eicosanoids may therefore play a pivotal role in daphnid reproduction. A wealth of synthetic and natural chemicals may affect invertebrate reproduction through endocrine disruption with one of the best known examples being imposex (masculinisation) of female molluscs caused by exposure to tributyl tin (TBT) [13]. It is therefore important to understand eicosanoid biosynthesis in *Daphnia*, and invertebrates in general, to fully recognize the potential mode of action of endocrine disrupters and how they may affect natural invertebrate populations.

Recently, the genome of *D. pulex* was fully sequenced [14], and action within the *Daphnia* Genomics Consortium has already been taken to start sequencing the *D. magna* genome [15,16]. In the meantime genes identified in *D. pulex* serve as a model for understanding eicosanoid biosynthesis, control and disruption in *Daphnia*. Here we present an overview of the putative eicosanoid biosynthesis in *Daphnia* based on annotation of genes from the *D. pulex* genome supported by recently published transcriptomic data (real-time quantitative PCR) of ibuprofen-stressed *D. magna* [12].

Methods

Putative genes related to eicosanoid biosynthesis were identified on the *D. pulex* genome website [14] through using several bioinformatic search tools such as GO (Gene Ontology) [17], KEGG (Kyoto Encyclopedia of Genes and Genomes) [18] and matches against InterPro protein domains. Annotation of these genes were verified through BLAST searches performed against Swissprot protein and non-redundant protein sequence databases via the *D. pulex* genome website [14].

Invertebrate and vertebrate COX protein sequences were retrieved from GenBank [19] and Ensembl [20] for phylogenetic analysis. Since no arthropod COX protein sequences were available that could aid the phylogeny with respect to *D. pulex*, we searched the GenBank Expressed Sequence Tag (EST) division (est_others) using

the BLAST algorithm (tblastn) to obtain additional arthropod COX sequences. Sequences that significantly resembled the sea squirt *Ciona intestinalis* COX amino acid sequence (E-value < 1e-20) were retrieved and included in the dataset. Six ESTs were obtained, representing two malacostracan species (Crustacea): Four ESTs were derived from *Homarus americanus* (accession numbers [DV772953](#), [DV774102](#), [EH401871](#) and [FD699680](#)) and two were derived from *Petrolisthes cinctipes* (accession numbers [FE773225](#), [FE820815](#)). After translating the nucleotide sequences to putative amino acids, it appeared that one *H. americanus* sequence ([FD699680](#)) did not overlap with the other three sequences. The three remaining sequences constituted two slightly deviating amino acid sequences; [DV774102](#) differed from [EH401871](#) and [DV772953](#). The four *H. americanus* sequences were combined to one sequence with variable and missing positions specified as 'unknown'. Furthermore, the inferred amino acid sequences of the two *P. cinctipes* ESTs were similar, and *P. cinctipes* was thus included once in the dataset. Note that these additional crustacean entries were EST derived, single pass sequenced, and therefore not guaranteed to be free from sequencing errors.

The sequence COX dataset was aligned using ClustalW2 [21], manually edited after inspection and subsequently analysed using the Gblocks web-server [22] to pinpoint and remove unreliably aligned regions. This analysis allowed for gaps in the final alignment. ModelGenerator [23] was applied to obtain a model of sequence evolution

(gamma distribution with four rate categories) using the Akaike Information Criterion. The predicted model, WAG+I+G, was specified in Phyml v2.4.4 [24] and a dendrogram was constructed using Maximum Likelihood (BIONJ [25] starting tree). Phylogenetic analysis used 100 bootstrap replicates. Finally, the obtained topology was visualised using TreeView [26].

Results and Discussion

Table 1 shows the putative genes related to eicosanoid biosynthesis that were identified through searching the *D. pulex* genome using several different bioinformatic tools (see Methods). Many of these genes had high similarity to their counterparts in higher organisms (Table 1). The bioinformatic evidence from *D. pulex* suggested that only the cyclooxygenase (COX) and lipoxygenase (LOX) pathways are present in *Daphnia* (Fig. 2) compared with the three pathways known from mammalian systems, i.e. the COX, LOX and epoxygenase pathways. This agrees with earlier findings as no epoxygenase pathway has been identified in invertebrates to date [1].

Both the COX and LOX pathways in *Daphnia* appeared to have a simpler structure than their mammalian counterparts (for comparison, see Figs. 1 and 2). For instance, there was no bioinformatic evidence of prostacyclin synthase, which converts PGG₂ into PGI₂, in the daphnid COX pathway. The gene encoding this enzyme was likewise not identified in the genome of the urochordate *C. intestinalis* [2]. Additionally, there was only bioinformatic

Table 1: Putative *Daphnia pulex* genes associated with eicosanoid biosynthesis

Putative gene ^a	Protein ID ^b	Protein size (AA)	Best match (species) ^c	Accession number	Similarity (%)	E-value
Phospholipase A2	DAPPUDRAFT_5959	129	<i>Apis mellifera</i>	XP_624621	76	0.0
Cyclooxygenase	DAPPUDRAFT_313427	689	<i>Sus scrofa</i>	NP_999486	62	0.0
Prostaglandin D2 synthase A	DAPPUDRAFT_307787	211	<i>Tribolium castaneum</i>	XP_970647	49	2.5E-35
Prostaglandin D2 synthase B	DAPPUDRAFT_316534	228	<i>T. castaneum</i>	XP_967406	54	2.3E-32
Prostaglandin E2 synthase	DAPPUDRAFT_56335	283	<i>Aedes aegypti</i>	EAT39424	61	0.0
Carbonyl reductase I (PG 9-ketoreductase)	DAPPUDRAFT_310758	257	<i>Xenopus laevis</i>	Q6DJN9	76	0.0
<i>Thromboxane A</i>	DAPPUDRAFT_96715	463	<i>Homarus americanus</i>	AAC28351	49	0.0
<i>Thromboxane B</i>	DAPPUDRAFT_328913	501	<i>H. americanus</i>	AAC28351	59	0.0
Lipoxygenase 1	DAPPUDRAFT_311736	562	<i>T. castaneum</i>	XP_969219	48	0.0
Lipoxygenase 2	DAPPUDRAFT_95367	433	<i>Bos taurus</i>	XP_593842	33	2.1E-43
Glutathione peroxidase	DAPPUDRAFT_337058	219	<i>Ixodes scapularis</i>	AAV66814	67	0.0
<i>Leukotriene A4 hydrolase</i>	DAPPUDRAFT_313156	619	<i>X. tropicalis</i>	NP_001006898	67	0.0
<i>Leukotriene B4 12-hydroxydehydrogenase</i>	DAPPUDRAFT_311788	340	<i>A. aegypti</i>	EAT35376	65	0.0
Prostanoid receptor EP4 isoform A	DAPPUDRAFT_58618	166	<i>Danio rerio</i>	LOC560057	59	3.6E-38
Prostanoid receptor EP4 isoform B	DAPPUDRAFT_58558	121	<i>Tetraodon nigroviridis</i>	CAF96390	60	4.2E-30

^aGene names displayed in italics are unlikely to be involved in daphnid eicosanoid biosynthesis (see Fig. 2 and text for further information); ^b*D. pulex* protein ID is linked to a GenBank [19] accession no.; ^cVertebrate species are signified by bold font, while invertebrate matches are denoted by normal font.




Figure 2

Putative eicosanoid biosynthesis pathway in *Daphnia* based on bioinformatic and transcriptomic evidence from *D. pulex* and *D. magna*. All the putative genes (names in italics) have been identified through different gene models in the *D. pulex* genome and are shown in black or grey font based on high (> 60%) or low (< 60%) similarity to proteins from other genomes (Table 1). Eicosanoids in grey font are less likely to be present in daphnids. Expression of ortholog genes in ibuprofen-stressed *D. magna* (24 h exposure to 20–80 mg l⁻¹) was analysed using real-time quantitative PCR [12]. Fold change difference in gene expression (mean ± SE) is shown relative to controls (grey values are only weakly significant). The enzyme LTB₄DH (encoded by *Ltb4dh*), that catabolises PGE₂, PGF_{2α} and LTB₄ to become inactive eicosanoids, is also known as 15-oxo-prostaglandin 13-reductase. All the specified eicosanoids have been identified in other arthropod species [1,2], except for the leukotrienes where only indirect evidence exists [37]. Abbreviations: HETE, hydroxyeicosatetraenoic; HPETE, hydroperoxyeicosatetraenoic acid. See text for further details.

evidence of one gene encoding COX in the *D. pulex* genome. A phylogenetic comparison of the predicted *D. pulex* COX with other protein sequences revealed that daphnid COX clusters with the invertebrates being most closely related to other crustaceans. The COX phylogeny likewise showed that COX-1 and COX-2 comprise two distinct clades amongst the vertebrates (Fig. 3). Generally, it is understood that invertebrates and lower vertebrates only have one non-specific type of COX [2], but recently two COX isoforms have been identified in the corals *Plexaura homomalla* [7] and *Gersemia fruticosa* [27]. Rowley et al. [2] suggest that the COX genes found in corals are an early version that predates the (supposed) vertebrate duplication into the typical constitutive COX-1 and inducible COX-2 isozymes found in vertebrates. Only one copy of the COX gene has been identified in the *C. intesti-*

nalis genome [2] which, as a member of the Phylum Chordata, shares a more recent common ancestor with the vertebrates. This was supported by our phylogenetic analysis (Fig. 3) suggesting that a duplication of the COX gene occurred in the Chordata. Rowley and colleagues further speculate that the evolution of COX-1 and COX-2 probably predates the emergence of bony fish some 350 million years ago [2]. This is supported by the fact that only one constitutively expressed COX isoform has been identified in the shark *Squalus acanthias* (Spiny dogfish) [2]. In our phylogenetic analysis *S. acanthias* COX clustered with the vertebrate COX-1 clade.

Like the putative *Daphnia* COX pathway, the structure of the daphnid LOX pathway also appeared to be more simple than its mammalian counterpart, which confirms pre-




Figure 3

Phylogenetic tree of cyclooxygenase (COX) based on protein sequences from a diverse range of organisms constructed using Maximum Likelihood. All bootstrap values above 80 have been removed. Scale: 0.1 substitutions per site. Sequences can be retrieved from GenBank [19] or Ensembl [20], while the *Daphnia pulex* COX was derived from a predicted protein sequence based on the best gene model available on the *D. pulex* genome portal [14]. Note that some sequences are based on EST information (see Methods). Abbreviations (accession no. in brackets with COX1 shown first when there are two no.): CANIS, *Canis familiaris* (NP_001003023; NP_001003354); CHICK, *Gallus gallus* (P27607); CIOIN, *Ciona intestinalis* (ENSCINT00000012734); CIOSA, *C. savignyi* (ENSCSAVT00000000782); DANRE, *Danio rerio* (Q8JH44; Q6P4V3); DPULX, *Daphnia pulex* (NCBI_GNO_0900053); FUNHE, *Fundulus heteroclitus* (Q6QNF2); GEFRU, *Gersemia fruticosa* (Q9GPF4; Q6S375); HOMA, *Homarus americanus* (EST: DV772953, DV774102, EH401871 and FD699680); HUMAN, *Homo sapiens* (P23219; P35354); MOUSE, *Mus musculus* (Q543T1; Q05769); ONCMY, *Oncorhynchus mykiss* (Q9DEQ0; Q9W715); PANTR, *Pan troglodytes* (ENSPTRT00000042391; ENSPTRT00000003246); PETRO, *Petrolisthes cinctipes* (EST: FE773225, FE820815); PLEHO, *Pleuraea homomalla* (Q96218; Q51X63); RABIT, *Oryctolagus cuniculus* (O97554; O02768); RAT, *Rattus norvegicus* (Q63921; Q63124); SALFO, *Salvelinus fontinalis* (Q9PTN3; Q9PW89); SHEEP, *Ovis aries* (P05979; P79208); SQUAC, *Squalus acanthias* (Q8UVQ3); XENLA, *Xenopus laevis* (A0A9J3); XENTR, *X. tropicalis* (ENSXETT00000035660; Q501R2).

vious findings [1]. There was gene sequence evidence that indicated the presence of three leukotrienes (LT) in *Daphnia*: LTA₄, LTB₄ and 12-oxo LTB₄ (Table 1). This is, however, doubtful since no LTs, nor the precursor 5-HPETE, have been identified in invertebrates and lower vertebrates to date [28]. It is possible that the putative LTA₄ hydrolase (LTA4H), which is a bi-functional enzyme in mammals, having both LTA₄ hydrolase and aminopeptidase activity, only has aminopeptidase activity in daphnids as has been reported in *Caenorhabditis elegans* [29]. However, recent evidence shows that bi-functionality of LTA4H is only six point mutations away in yeast compared to the mammalian enzyme [30]. There is reason to believe that daphnid LTA4H may be bi-functional, and thus able to convert LTA₄ into LTB₄; a hypothesis that we will test in future experiments. Furthermore, transcriptomic evidence from *D. magna* shows that the expression of *leukotriene B4 12-hydroxydehydrogenase* (*Ltb4dh*) increases with increasing concentration of the eicosanoid-inhibiting drug ibuprofen [11]. This does not prove that LTs are present in daphnids, but yet again we cannot rule out the possibility. The enzyme (LTB4DH) encoded by *Ltb4dh* is also bi-functional in mammals regulating eicosanoids by rapidly degrading different E and F series PGs and LTB₄ [31] (see below). The former function of LTB4DH (PG catabolism) appears to be the most likely in daphnids until solid proof exists about the presence of LTs (Fig. 2).

The presence of lipoxins is, however, more likely, and bioinformatic information from *D. pulex* suggests that at least two LOX enzymes are present (Table 1). LOX enzymes have been found in all organisms studied, from bacteria to man. The two putative *D. pulex* LOXs are both composed by two domains; an N-terminal lipase domain belonging to the InterPro protein family 734 (IPR000734) and a C-terminal LOX LH2 domain (IPR001024). This resembles mammalian LOX enzymes that are also comprised of two domains; a regulatory N-terminal domain that is similar to mammalian lipases and a catalytic LOX domain (C-terminal) [32]. LOX LH2 is the only LOX related domain that has been identified in the *D. pulex* genome. Other known LOX domains include e.g. mammalian LOX (IPR001885) and LOX, C-terminal (IPR013819). There are 13 *D. pulex* gene models that contain the LOX LH2 domain [14], but only two genes contain both an N-terminal domain similar to mammalian lipases (IPR000724) and a C-terminal LOX domain (Table 1). Further investigations are needed to specify what type of LOX enzyme these two *Daphnia* sequences represent prior to analysing their phylogenetic relationship. However, it is likely that they could be 8-LOX and 12-LOX, which synthesise different lipoxins from 8-HPETE and 12-HPETE, as these LOXs have been identified in a range of invertebrate species [8,33,34].

Two enzymes, 15-hydroxyprostaglandin dehydrogenase (PGDH) and LTB4DH are known to irreversibly inactivate bioactive eicosanoids in mammals. Both enzymes are key in regulating the hormonal-like action of eicosanoids by rapidly degrading PGE₂, PGF_{2α} and LTB₄, as overproduction of these potent mediators may have serious physiological effects such as initiating inflammation [31]. It appeared that LTB4DH fulfils this regulatory function single-handedly in daphnids, as there was no indication of PGDH being present (Table 1, Fig. 2).

The bioinformatic and transcriptomic evidence from *D. pulex* and *D. magna* (Fig. 2) suggests that PGs (e.g. PGH₂, PGE₂ and PGF_{2α}), lipoxins and possibly LTs could be present in daphnids. Low similarity of TXA₂ synthase and PGD₂ synthase to ortholog proteins from other genomes (Fig. 2) render the presence of PGD₂ and TXA₂ to be less certain in daphnids, although this could merely be due to daphnids having more divergent versions of these proteins. Nevertheless, it seems most likely that daphnids do not produce TXA₂ since the gene encoding TXA₂ synthase has not been identified in the genome of *C. intestinalis* [2]. The LOX encoding genes identified in *D. pulex* were also slightly doubtful due to the same reasons, but it would be more probable that these enzymes are present in *Daphnia* as both 8-LOX and 12-LOX derived lipoxins are common in invertebrates [8,33,34]. PGA₂ may also be present in daphnids as it is non-enzymatically rearranged from PGE₂ and has been detected in several arthropods [1]; but until verified by mass spectrometry or the like it remains speculative what eicosanoids are present in *Daphnia*. Moreover, the annotation of genes from the daphnid eicosanoid biosynthesis (and other daphnid genes for that matter) should improve as more invertebrate genomes become sequenced and annotated.

The possible roles of eicosanoids in daphnids have already been suggested from several invertebrate studies, including *D. magna* [12], where both prostanoids and lipoxigenase products appear to be important agents in oogenesis (especially vitellogenesis) and embryogenesis [35-38]. For instance, PGE₂ is known to initiate egg-laying behaviour in several insect species (e.g. orthopterans), where it seems to regulate muscle contractions in the ovarian musculature [39]. Furthermore, many of the above-mentioned eicosanoids have likewise been identified as important mediators in arthropod immune systems (both COX and LOX products) [40] and ion transport physiology (mainly PGE₂ and PGF_{2α}) [1]. Until more integrated phenotypic and genomic evidence exists it is difficult to infer an exact role for eicosanoids in daphnids, as they may be involved in several processes and act in different tissues. Nevertheless, it is almost certain that eicosanoids play vital roles in the functioning of processes key to daphnid reproduction and survival [9,10,12]. Finally, bio-

informatic evidence from the *D. pulex* genome also revealed that two prostanoid G-protein coupled receptors may be present, thus further supporting the evidence that eicosanoids are bioactive agents in daphnids (Table 1).

Conclusion

Eicosanoids are key molecules, involved in the function of fundamentally important biological systems. A better understanding of their biochemistry and genetic control in invertebrates will help to improve our understanding of their significance in these organisms. Here we have outlined a putative structure of eicosanoid biosynthesis in *Daphnia*, a key macroinvertebrate in freshwater ecosystems. It would seem, from transcriptomic and phenotypic evidence, that eicosanoids play a pivotal role in daphnid reproduction [12]; but their importance in other physiological functions such as the immune system remains to be investigated. Improved knowledge of the function and synthesis of eicosanoids in *Daphnia* and other invertebrates could have very important implications for several areas within ecology including ecological risk assessment. This provisional overview of daphnid eicosanoid biosynthesis provides a guide on where to focus future research activities in this area.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LHH and MJTNT performed the bioinformatic and phylogenetic analyses. LHH drafted the manuscript. All authors contributed intellectually to the manuscript and approved the final version.

Acknowledgements

We gratefully acknowledge the financial support of AstraZeneca, Syngenta, NERC (project NER/D/S/2002/00413 "The population and molecular stress responses of an ecotoxicology indicator species") and The Research Endowment Trust Fund of the University of Reading. We thank the *Daphnia* Genomics Consortium (DGC) for giving us the opportunity of annotating *D. pulex* genes related to eicosanoid biosynthesis, and Vibeke Simonsen (University of Aarhus, National Environmental Research Institute) for critically reading the manuscript. The sequencing and portions of the analyses were performed at the DOE Joint Genome Institute under the auspices of the U.S. Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48, Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231, Los Alamos National Laboratory under Contract No. W-7405-ENG-36 and in collaboration with the DGC [16]. Additional analyses were performed by wFleaBase [15], developed at the Genome Informatics Lab of Indiana University with support to Don Gilbert from the National Science Foundation and the National Institutes of Health. Coordination infrastructure for the DGC is provided by The Center for Genomics and Bioinformatics at Indiana University, which is supported in part by the MET-ACyt Initiative of Indiana University, funded in part through a major grant from the Lilly Endowment, Inc. Our work benefits from, and contributes to the DGC [16].

References

1. Stanley DW: **Eicosanoids in invertebrate signal transduction systems.** New Jersey, Princeton University Press; 2000.
2. Rowley AF, Vogan CL, Taylor GW, Clare AS: **Prostaglandins in non-insectan invertebrates: recent insights and unsolved problems.** *Journal of Experimental Biology* 2005, **208**(1):3-14.
3. McGettigan P, Henry D: **Current problems with non-specific COX inhibitors.** *Current Pharmaceutical Design* 2000, **6**(17):1693-1724.
4. Ye XQ, Hama K, Contos JJA, Anliker B, Inoue A, Skinner MK, Suzuki H, Amano T, Kennedy G, Arai H, Aoki J, Chun J: **LPA(3)-mediated lysophosphatidic acid signalling in embryo implantation and spacing.** *Nature* 2005, **435**(7038):104-108.
5. Funk CD: **Prostaglandins and leukotrienes: Advances in eicosanoid biology.** *Science* 2001, **294**(5548):1871-1875.
6. Feige JN, Gelman L, Michalik L, Desvergne B, Wahli W: **From molecular action to physiological outputs: Peroxisome proliferator-activated receptors are nuclear receptors at the crossroads of key cellular functions.** *Progress in Lipid Research* 2006, **45**(2):120-159.
7. Valmsen K, Jarving I, Boegliin WE, Varvas K, Koljak R, Pehk T, Brash AR, Samel N: **The origin of 15R-prostaglandins in the Caribbean coral *Plexaura homomalla*: Molecular cloning and expression of a novel cyclooxygenase.** *Proceedings of the National Academy of Sciences of the United States of America* 2001, **98**(14):7700-7705.
8. Ragab A, Durand J, Bitsch C, Chap H, Rigaud M: **The lipoxygenase pathway of arachidonic acid metabolism in reproductive tissues of the firebrat, *Thermobia domestica* (Thysanura).** *Insect Biochemistry* 1991, **21**(3):321-326.
9. Hayashi Y, Heckmann LH, Callaghan A, Sibly RM: **Reproduction recovery of the crustacean *Daphnia magna* following chronic exposure to ibuprofen.** *Ecotoxicology* 2008, **17**:246-251.
10. Heckmann LH, Callaghan A, Hooper HL, Connon R, Hutchinson TH, Maund SJ, Sibly RM: **Chronic toxicity of ibuprofen to *Daphnia magna*: Effects on life history traits and population dynamics.** *Toxicology Letters* 2007, **172**(3):137-145.
11. Heckmann LH, Connon R, Hutchinson TH, Maund SJ, Sibly RM, Callaghan A: **Expression of target and reference genes in *Daphnia magna* exposed to ibuprofen.** *BMC Genomics* 2006, **7**:175.
12. Heckmann LH, Sibly RM, Connon R, Hooper HL, Hutchinson TH, Maund SJ, Hill CJ, Bouetard A, Callaghan A: **Systems biology meets stress ecology: Linking molecular and organismal stress responses in *Daphnia magna*.** *Genome Biology* 2008, **9**(2):R40.
13. Hutchinson TH: **Reproductive and developmental effects of endocrine disrupters in invertebrates: in vitro and in vivo approaches.** *Toxicology Letters* 2002, **131**(1-2):75-81.
14. **JGI Genome Portal** [<http://www.jgi.doe.gov/Daphnia>]
15. **wFleaBase** [<http://wFleaBase.org>]
16. ***Daphnia* Genomics Consortium** [<http://daphnia.cgb.indiana.edu>]
17. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G: **Gene Ontology: tool for the unification of biology.** *Nature Genetics* 2000, **25**(1):25-29.
18. Kanehisa M, Goto S: **KEGG: Kyoto Encyclopedia of Genes and Genomes.** *Nucleic Acids Research* 2000, **28**(1):27-30.
19. **GenBank** [<http://www.ncbi.nlm.nih.gov/Genbank/>]
20. **Ensembl** [<http://www.ensembl.org/>]
21. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD: **Multiple sequence alignment with the Clustal series of programs.** *Nucleic Acids Research* 2003, **31**(13):3497-3500.
22. Castresana J: **Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis.** *Mol Biol Evol* 2000, **17**(4):540-552.
23. Keane TM, Creevey CJ, Pentony MM, Naughton TJ, McLnerney JO: **Assessment of methods for amino acid matrix selection and their use on empirical data shows that ad hoc assumptions for choice of matrix are not justified.** *BMC Evolutionary Biology* 2006, **6**:29.
24. Guindon S, Gascuel O: **A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood.** *Systematic Biology* 2003, **52**(5):696-704.

25. Gascuel O: **BIONJ: An improved version of the NJ algorithm based on a simple model of sequence data.** *Molecular Biology and Evolution* 1997, **14**:685-695.
26. Page RDM: **TreeView: an application to display phylogenetic trees on personal computers.** *Comput Appl Biosci* 1996, **12(4)**:357-358.
27. Järving R, Järving I, Kurg R, Brash AR, Samel N: **On the evolutionary origin of cyclooxygenase (COX) isozymes - Characterization of marine invertebrate COX genes points to independent duplication events in vertebrate and invertebrate lineages.** *Journal of Biological Chemistry* 2004, **279(14)**:13624-13633.
28. Morgan EL, Maskrey BH, Rowley AF: **At what stage in metazoan evolution did leukotriene generation first appear? - key insights from cartilaginous fish.** *Developmental and Comparative Immunology* 2005, **29(1)**:53-59.
29. Baset HA, Ford-Hutchinson AW, O'Neill GP: **Molecular cloning and functional expression of a *Caenorhabditis elegans* aminopeptidase structurally related to mammalian leukotriene A(4) hydrolases.** *Journal of Biological Chemistry* 1998, **273(43)**:27978-27987.
30. Tholander F, Kull F, Ohlson E, Shafqat J, Thunnissen MMGM, Haeggström JZ: **Leukotriene A4 hydrolase, insights into the molecular evolution by homology modeling and mutational analysis of enzyme from *Saccharomyces cerevisiae*.** *J Biol Chem* 2004, **280(39)**:33477-33486.
31. Hori T, Yokomizo T, Ago H, Sugahara M, Ueno G, Yamamoto M, Kumasaka T, Shimizu T, Miyano M: **Structural basis of leukotriene B-4 12-hydroxydehydrogenase/15-oxo-prostaglandin 13-reductase catalytic mechanism and a possible Src homology 3 domain binding loop.** *Journal of Biological Chemistry* 2004, **279(21)**:22615-22623.
32. Gillmor SA, Villaseñor A, Fletterick R, Sigal E, Browner MF: **The structure of mammalian 15-lipoxygenase reveals similarity to the lipases and the determinants of substrate specificity.** *Nature Structural Biology* 1997, **4(12)**:1003-1009.
33. Knight J, Taylor GW, Wright P, Clare AS, Rowley AF: **Eicosanoid biosynthesis in an advanced deuterostomate invertebrate, the sea squirt (*Ciona intestinalis*).** *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids* 1999, **1436(3)**:467-478.
34. Kuo JM, Pan BS, Zhang HJ, German JB: **Identification of 12-Lipoxygenase in the hemolymph of tiger shrimp (*Penaeus japonicus* Bate).** *Journal of Agricultural and Food Chemistry* 1994, **42(8)**:1620-1623.
35. Medeiros MN, Mendonca LH, Hunter AL, Paiva-Silva GO, Mello FG, Henze IP, Masuda H, Maya-Monteiro CM, Machado EA: **The role of lipoxygenase products on the endocytosis of yolk proteins in insects: Participation of cAMP.** *Archives of Insect Biochemistry and Physiology* 2004, **55(4)**:178-187.
36. Sagi A, Silkovsky J, Fleisher-Berkovich S, Danon A, Chayoth R: **Prostaglandin E(2) in previtellogenic ovaries of the prawn *Macrobrachium rosenbergii*: Synthesis and effect on the level of cAMP.** *General and Comparative Endocrinology* 1995, **100(3)**:308-313.
37. Silver RB, Oblak JB, Jeun GS, Sung JJ, Dutta TC: **Leukotriene B4, an arachidonic acid metabolite, regulates intracellular free calcium release in eggs and mitotic cells of the sand dollar (*Echinarracnius parma*).** *Biological Bulletin* 1994, **187(2)**:242-244.
38. Spaziani EP, Hinsch GW, Edwards SC: **Changes in prostaglandin E(2) and F2-alpha during vitellogenesis in the Florida crayfish *Procambarus paeninsulanus*.** *Journal of Comparative Physiology B: Biochemical, Systemic and Environmental Physiology* 1993, **163(7)**:541-545.
39. Loher W, Ganjian I, Kubo I, Stanley-Samuels D, Tobe SS: **Prostaglandins - Their role in egg-laying of the cricket *Teleogryllus commodus*.** *Proceedings of the National Academy of Sciences of the United States of America* 1981, **78(12)**:7835-7838.
40. Miller JS, Nguyen T, Stanley-Samuels DW: **Eicosanoids mediate insect nodulation responses to bacterial infections.** *Proceedings of the National Academy of Sciences of the United States of America* 1994, **91(26)**:12418-12422.
41. **KEGG: Kyoto Encyclopedia of Genes and Genomes** [<http://www.genome.jp/kegg/>]

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

